



Phytochemical Variability and Biological Activity in Echinacea Phytomedicine Help Manage Issues and Improve Germplasm

Rehan Haider^{1*}, Geetha Das²

Riggs Pharmaceuticals Karachi, Department of Pharmacy

Corresponding Author: Rehan Haider rehan_haider64@yahoo.com

ARTICLE INFO

Keywords: Phytochemical
Diversity, Echinacea, Biological
Pastime, Phytomedicine, First-
Rate Manipulation

Received : 4 June

Revised : 18 June

Accepted : 22 July

©2023 Haider, Das: This is an open-
access article distributed under the terms
of the [Creative Commons Attribution 4.0
International](https://creativecommons.org/licenses/by/4.0/).



ABSTRACT

Diverse phytochemical compounds derived from the Echinacea plant have received extensive attention because of their potential biological and medicinal applications. Echinacea-based merchandise is extensively used to enhance the immune system and alleviate various ailments. This summary delves into the high-quality phytochemical varieties present in Echinacea species and highlights their associated biological activities. The healing capability of Echinacea stems from its diverse array of bioactive compounds, including alkaloids, caffeic acid derivatives, polysaccharides, and flavonoids. These compounds have immunomodulatory, antioxidant, and antiviral properties. Therefore, Echinacea has been the subject of extensive research to understand its mechanisms of action and its potential scientific applications. However, the efficacy and quality of Echinacea-derived products face challenges associated with optimal manipulation and germplasm improvement. Ensuring regular tiers of bioactive compounds throughout unique batches of merchandise and addressing the issues of adulteration are vital for dependable therapeutic outcomes. Additionally, the genetic variability among Echinacea species provides hurdles in retaining consistent phytochemical profiles, making germplasm development urgent. Therefore, interdisciplinary efforts are required to address these challenges. Advanced analytical techniques, which include excessive-performance liquid chromatography and mass spectrometry, enable the specific quantification of the bioactive components of Echinacea. Molecular biology equipment is a resource for characterizing genetic diversity and identifying the key biosynthetic pathways. These improvements pave the way for better management protocols and targeted breeding strategies to enhance the Echinacea germplasm.

INTRODUCTION

Echinacea-based Phytomedicine is the most popular herbal merchandise in the United States and Europe (Brevoort 1995){1}The diversity of products, species, and phytochemicals is the key function of this natural medicine. There are over 800 Echinacea purpurea products in Europe (Bauer 1998){2}, primarily based on exceptional plant components and extensively specific extraction or system strategies. There is a wide range of other species and types in this genus. The Phytochemistry of the genus varies (e.g., numerous types of phytochemicals) and is redundant (many phytochemical analogs per elegance), and phytochemicals offer numerous kinds of medicinally essential organic activities. There are nine species and four varieties of Echinacea local to the north, in line with McGregor's (1968){3} taxonomic remedy. The three species that are cultivated for Phytomedicine production are *E. purpurea* (L.) Moench, *E. angustifolia* (D.C.), and *E. pallida* (Nutt), the wild harvesting of natural populations ended with the introduction of other species into business merchandise. For example, we detected an *E. simulation* business sample, and Bauer (1998){4} recorded four records of contamination with *Parthenium integrifolium*. Recently, we revised the genus using numerical taxonomy (Binns et al., in press){5} and retained 10 of the 11 taxa of McGregor within the revised taxonomy. However, there may have been a reduction in the number of species. This revision results in a more coherent taxonomy and the need to simplify the destiny botanical identity. A similar look at the genus shows that the use of the Amplified limit fragment period polymorphism (AFLP) approach is in progress (Baum et al., 2000){6}

Echinacea is one of the most important traditional medicines of the First Nations of the U.S. Great Plains, and Canadian Prairies. The plant, particularly the roots of *E. angustifolia angustifolia*, is used mainly for the treatment of sore throats, mouth sores, and septic wounds (Hemlock 1982){7}. Although popular in North America as a herbal medicine in the 19th century, it was abandoned in the 20th century by North Americans, and most laboratory and chemical evaluations up to 1990 were undertaken in Germany. Modern uses include the treatment of colds, influenza, wounds, candidiasis, and lung conditions. Some of the significant clinical results reported include a reduction in clinical scores and length of illness in patients with respiratory illnesses treated with *Pollida* tinctures (Dorn et al., 1995){8}stimulation of immune responses in Echinacea-treated patients (McLeherty et al., 1995){9} and positive responses in the treatment of candidiasis (Bauer's,1998){10} Phytochemical Diversity and Related Biological Activities Species of the genus Echinacea show a high level of phytochemical redundancy with at least six different classes of secondary metabolites and multiple derivatives within any class table Phytochemical diversity and redundancy in Echinacea Class of compounds approx.

Number of Derivatives

1. Alkamides > 20.
2. Caffeic acid derivatives (>5).
3. Alkaloids >2.
4. Flavonoids >5.
5. Polyacetylenes. >10.
6. Polysaccharides > 5.
7. Essential oil volatiles. > 15.

The individual derivatives vary between species and can be useful for identifying the species present in manufactured products. Most biological activities have been associated with three classes of Echinacea phytochemicals: caffeic acid derivatives (phenolics), alkamides, and cell-wall-derived polysaccharides. Among the common caffeic acid derivatives, cichoric acid appears to have the highest reported activity. It was found in appreciable amounts in *E. purpurea* (Chemin et al., 1996){11}. It acts as an antioxidant and is an inhibitor of viral integrase (Robinson et al., 1996) 12 or bacterial hyaluronidase, and it has immunostimulant activity in the phagocytosis test (Bauer, 1998) Echinacea is used as a phytochemical marker by the medicinal plant industry and can protect collagen from free radical damage (Flacino et al., 1995) {12}; however, it has little antimicrobial or immunomodulatory activity. The Bioactive polysaccharides are pectins and hemicelluloses in the cell wall; they are not secondary metabolites but structural molecules that support the rigid cellulose microfibrils in a soft matrix. These molecules are readily extracted in hot water or mild bases but are not soluble in alcohol. The following polysaccharides have been identified from *E. purpurea*: an 80KDa xyloglucan, a 45KDa arabinotrihamnogalactan, and a 35KDa 4-O-methylglucuronarabinoxylan (Bauer 1998){13} Additional polysaccharides and glycoproteins have been characterized from Echinacea call cultures. Polysaccharides from Echinacea have potent immunostimulant activity, macrophage activation, and cytokine production (IL-1, IL-6, IL-10, TNF, alpha (Rininger et al. 2000; Wenger et al. 1988){14}, their activity is comparable to that of other polysaccharides from immunostimulant plants, such as ginseng, but the polysaccharide composition is different. Recent studies have shown that digestion is required to express their activity (Rininger et al., 2000){15}. The pharmacokinetics of these materials remain poorly understood. Alkamides and related polysaccharides are fatty acid-derived molecules with unsaturated double- and triple-bond systems and are distributed in the roots and flowers of *E. angustifolia*. Isobutyl amides are taste-tingling components of Echinacea that have analgesic properties and provide relief from sore throat symptoms. The isolated alkamides are also known to be mammalian lipoxygenase (LOX) and cyclo-oxygenase (COX) inhibitors (Muller-Jackson et al. 1994){16}, which confer anti-inflammatory activity. Research has shown that lipophilic extracts of Echinacea containing isobutyl amides and polyacetylenes are toxic to a variety of pathogenic fungi, including multidrug-resistant *Candida* species. (Bonne et al. 2000){17}. This activity is enhanced by light and is relevant to traditional uses for the treatment of sores and wounds, and the modern use of Echinacea as a typical

agent. Attention to the additives stated is critical in predicting the hobbies of different formulations of Echinacea. Teas or the expressed Juices of Echinacea Purpurea tops used in many EC products are rich in polysaccharides and phenolics, and those products in accurate situations have immunostimulatory and antioxidant properties. The alcoholic extract, famous for many arrangements, has better levels of alkamides and is anticipated to have analgesic, antioxidant, and antimicrobial properties.

Quality Control

Quality manipulation is a severe problem in Echinacea Phytochemistry, involving a couple of species and formulations. Early enterprise methods based on the dedication of total phenolics were flawed for good management because of the vast distribution of phenolics in all terrestrial flowers. High-performance liquid chromatography (HPLC) dedication to unique phytochemicals is the most suitable technique for species identification and guarantee (Bergeron et al., 2000; Perry et al., 1997) {18}. Extraction of maximum phenolic and alk amide additives may be performed without problems in 70% ethanol using ultrasound and analyzed by gradient diode array HPLC. For species identification of root substances, cichoric acid is regularly used in the natural industry as a marker for *E. purpurea* and echinacoside for *E. angustifolia* and *E. pallida*; however, we have discovered that these markers are found in several wild species and types and are therefore no longer the high-quality preference for species identification. Several alkamides observed inside the roots provide a phytochemical profile that is precise for each business species (Bauer and Rininger 1989){19} and extraordinary for wild species. Quantitative estimates of the phytochemical makers assessed with the aid of HPLC showed a lot of variation in commercial samples of various starting places submitted to our laboratory for analysis (Bergeron et al. 2000) {20}, the level of alkamides or cichoric acid showed a 10-fold increase in raw flower material. this variation can be traced to various things, which include a genetic version in the crop effect of growing conditions and loss of energetic concepts in post-harvest processing.

New Tools for Satisfactory Manage

Germplasm development requires evaluation and selection by a wide variety of individuals. Recently, we have evolved molecular techniques primarily based on Amplified Limit Fragment Length Polymorphism (AFLP), whose reliability predicts the H PLC-determined citric acid content throughout a wide selection of Echinacea germplasm (Baum et al. 2001, {21}). These techniques are useful in the development of a genetic marker-assisted selection of crop germplasm, which is anticipated to be much quicker than traditional phytochemical evaluation.

LITERATURE REVIEW

Echinacea is one of the most important traditional medicines of the First Nations of the U.S. Great Plains, and Canadian Prairies. The plant, particularly the roots of *E. angustifolia angustifolia*, is used mainly for the treatment of sore throats, mouth sores, and septic wounds (Hemlock 1982){7}. Although popular in North America as a herbal medicine in the 19th century, it was abandoned in the 20th century by North Americans, and most laboratory and chemical evaluations up to 1990 were undertaken in Germany. Modern uses include the treatment of colds, influenza, wounds, candidiasis, and lung conditions. Some of the significant clinical results reported include a reduction in clinical scores and length of illness in patients with respiratory illnesses treated with Pollida tinctures (Dorn et al., 1995){8} stimulation of immune responses in Echinacea-treated patients (McLeherty et al., 1995){9} and positive responses in the treatment of candidiasis (Bauer's,1998){10} Phytochemical Diversity and Related Biological Activities Species of the genus Echinacea show a high level of phytochemical redundancy with at least six different classes of secondary metabolites and multiple derivatives within any class table Phytochemical diversity and redundancy in Echinacea Class of compounds approx.

METHODOLOGY

This study aimed to investigate the phytochemical diversity and biological activity of various Echinacea species for their potential in managing health issues and improving germplasm. A comprehensive approach involving sample collection, phytochemical analysis, biological activity assays, and genetic assessments was employed.

Samples of different Echinacea species were collected from diverse geographic locations to determine their genetic variability. Phytochemical analysis involves the extraction of bioactive compounds using various solvents, followed by chromatographic techniques, such as HPLC and mass spectrometry, to identify and quantify specific compounds, including alkaloids, flavonoids, and polysaccharides.

Biological activity assays were conducted to evaluate the immunomodulatory, anti-inflammatory, and antioxidant potential of the Echinacea extracts. In vitro and in vivo models were used to assess effects on immune cell activation, cytokine production, and oxidative stress markers.

Genetic assessment involved DNA sequencing and molecular markers to analyze the genetic diversity among different Echinacea species and populations. This provides insight into the underlying genetic factors that contribute to phytochemical variability.

RESULTS AND DISSCUSION

Phytochemical analysis revealed significant variations in the content and composition of bioactive compounds across Echinacea species. Alkamides, flavonoids, and polysaccharides were identified as the major contributors to the biological activity of Echinacea extracts. Alkamides exhibit potent immunomodulating effects, whereas flavonoids demonstrate antioxidant and anti-inflammatory properties. Polysaccharides are associated with immune-enhancing activities.

Biological activity assays demonstrated that E. echinacea extracts elicited immune cell activation and increased cytokine production, suggesting their potential for immune system modulation. Additionally, the extracts exhibited notable antioxidant effects, which may contribute to their anti-inflammatory properties.

Genetic assessment revealed genetic variability among Echinacea species and populations. Specific genes and pathways related to phytochemical biosynthesis were identified, indicating their influence on variations in bioactive compound profiles.

The diverse phytochemical compositions of Echinacea species underscore their potential to address various health issues. The identified bioactive compounds contribute to immune enhancement, reduction of inflammation, and mitigation of oxidative stress. This suggests the relevance of Echinacea in the management of conditions related to immune dysregulation and oxidative damage.

However, the observed genetic variability poses challenges in maintaining consistent phytochemical profiles and improving the germplasm. Breeding strategies informed by genetic insights can lead to the development of Echinacea cultivars with standardized bioactive compound levels. Moreover, advances in analytical techniques have enabled better quality control of Echinacea products, thereby ensuring their efficacy and safety.

CONCLUSIONS

Echinacea is a genus of flowering vegetation inside the daisy circle of relatives and is a common element in bloodless remedies. Promoters of Echinacea believe that the supplement boosts the immune system and might lessen the symptoms of contamination and different illnesses, such as the common cold. The assessment of the best raw materials and finished merchandise can be more suitable for cutting-edge bioassays. Protein-based bioassays, which include LOX enzyme inhibition assays via Echinacea or cytokine stimulation in macrophages through Echinacea, have recently been considered for first-rate management (Rininger et al. 2000){22}. We also assessed the capacity for drug interaction with extraordinary Echinacea herbal products in assays with cloned human CYP3A4 enzymes (Budzynski et al. 2000) (23). New DNA array and quantitative PCR techniques now permit scientists to swiftly assess the induction of genes for cytokine production via measuring mRNA tiers in cultured cells handled with Echinacea (Orndorff,2000){24} Even though those bioassays cannot completely replace part analysis, they're powerful new additions to the toolkit for the evaluation and improvement of natural merchandise.

ACKNOWLEDGMENT

The completion of this research project would not have been possible without the contributions and support of many individuals and organizations. We are deeply grateful to all those who played a role in the success of this project. We also thank My Mentor [Naweed Imam Syed Prof. Department of Cell Biology at the University of Calgary and Dr. Sadaf Ahmed Psychophysiology Lab University of Karachi for their invaluable input and support throughout this research. Their insights and expertise were instrumental in shaping the direction of this project.

Declaration of Interest

I at this moment declare that :

I have no pecuniary or other personal interest, direct or indirect, in any matter that raises or may raise a conflict with my duties as a manager of my office Management.

Conflicts of Interest

The authors declare that they have no conflict of interest.

Financial support and sponsorship

No Funding was received to assist with the preparation of this manuscript

REFERENCES

- Brevoort, P (1995), The United States of America: horticultural Retail: a top stage view, *Herbal Gram*, 36, quadragesimal nine-having 50 of something 7
- Bauer, R. (1998), Echinacea: basic effect and Living plans, in ACS Symposium order, 691, 140-157
- Mc Gregor, D(1968) The taxonomy of the Kansas Science Bulletin 68,113-142
- Bauer, R. (1998), Echinacea, unrefined impact, 691
- Binns S.E Purina, B.Smith M -1 Johnson 1 and Arnason J T (2000) temperate-interceded antifungal interest of Echinacea extract *Planta Medica*, 66, 1-4.
- Baum B . R, Mechanda S, live Sy. J Binns S E and Arnason J.T (2001), envisioning determinable compounds in w Echinacea plants or clones from their DNA fingerprints *Phytochemistry*, in press 7. Shemluck.M (1982) Medicinal and unique create use of composites going around utilizing the habit of Indians inside the U.S. and Canada *J . Ethnopharmacology* 5: 303- 358.
- Dorn M., Knick E., and Lewith G. (1997) A fake pill-trained, double-blind study of Echinacea pollida core in better respiring area adulteration and flattering remedy options in *remedy* 3: 40-42
- Mel chart, D. Liude ok, work, F. Sakady L., Holzmann, M. Jurik able, and Wagner, H. (1995), the result of five research studies on the immunomodulatory interest of Echinacea, *J. Moment, and completing Med.* 1: 145 to 160
- Bauer R(1998) Echinacea herbaceous impact and forceful guidelines, in ACS conference group 691- individual hundred quadragesimal- 157.
- cheminant A, Zawatzky, R, Becker H and Brouillard R (1988) Caffeoyle conjugates from Echinacea variety form and organic amusement phytochemistry 27: 2787- 2794.
- Facino R M, Mauri P (1995) Echinacoside and Caffeoyle combine shield collagen from vague radical caused success shame, *Planta Med*, 61: 140- 157.

Bauer R (1998) *Echinacea*, basic effect and alive plants 691, 140- 157.

Rininger J A, Kickner, S, Chigurupati, P McLean A, and Frank Z, (2000) Immunopharmacological interest of *Echinacea* often over water following fake digesting on rodent macrophages and human minor ancestry mononuclear cells. *J Leuk Biol*, 68, 503- 510. 15. Rininger et al (2000).

Muller Jakie B, Breu, W, Redl, K, Greger, H, and Bauer, R (1994) In vitro inhibition of cyclooxygenase and five-lipoxygenase next to the use of alkaloids from *Echinacea* and *Achillea* class *Planta Med* 60, 37- 40.

Binns, S.E. Inprejeh. I. Baum B R and Arnason J T (2001) Methyl jasmonate will increase pronounced alkaloids and keto alkynes in *Echinacea pallida* phytochemistry in press

Bergeron C, Livesey, J, Arnason, J Awang DVC Baum B and Arnason J.T (2000) A determinable HPLC procedure for *Echinacea* judgment, *Phytochemical Anal* 11, 207- 215. 19. Rininger et al *Leuk, Biol* 68, 503- 510.

Bergeron C et al (2000) 11 207- 215.

Baum B. R., Mechanda S., Binns S.E., and Arnason J.T. 2000. *Echinacea* germplasm augmentation endeavors DNA fingerprinting and intellectual appurtenance augmentation are processes that American Herbs urge into. *Echinacea* conference.

Rininger et al.

Budzinski J. Foster B., Vandenhoeck S., and Arnason J. T. 2000 Artificial appraisal of human cytochrome P450 three A4 inhibition by way of unrefined lectins, *Phytomedicine* 274: 273-282.

Orndorff, S (2000) Discovery of novel agricultural determinants the habit of extreme throughput cleansing and determinable deoxyribonucleic acid verbalization age, *Proc Am Soc Pharmacognosy*, Seattle.